11. (Currently Amended) A bathophenanthroline compound of formula (I):

E

wherein R<sup>1</sup> and R<sup>2</sup> and are derived from R<sup>1</sup>-Li and R<sup>2</sup>-Li respectively, and may be the same or different and independently represent a hydrocarbon group provided that at least one of R<sup>1</sup> and R<sup>2</sup> has at least two carbons; and wherein R<sup>1</sup> and R<sup>2</sup> are selected from the group consisting of an ethyl group, an n-propyl group, an isopropyl group, an n-pentyl group, an iso-pentyl group, a neopentyl group, a tert-pentyl group, a cyclopentyl group, a methycyclopentyl methylcyclopentyl group, a dimethycyclopentyl dimethylcyclopentyl group, a trimethycyclopentyl trimethylcyclopentyl group, a tetramethycyclopentyl tetramethylcyclopentyl group, an n-hexyl group, a 2-ethylbutyl group, a 3,3-dimethylbutyl group, a cyclohexyl group, an n-cyclohexylmethyl group, an n,n-dimethycyclohexyl n,n-dimethylcyclohexyl group, an n,n,ntrimethycyclohexyl an n,n,n-trimethylcyclohexyl group, a tert-octyl group, a 2-ethylheyxl group, an n-nonyl group, an n-decyl group, an n-dodecyl group, an n-tetradecyl group, an n-hexadecyl group, a benzyl group, a phenethyl group, an  $\alpha$ -methylbenzyl group, an  $\alpha$ , a dimethylbenzyl group, a 1-naphthylmethyl group, a 2-naphthylmethyl group, a furfuryl group, a 2-methylbenzyl group, a 3-methylbenzyl group, a 4-methylbenzyl group, a 4-ethylbenzyl group, a 4isopropylbnezył 4-isopropylbenzył group, a 4-tert-butylbenzyl group, a 4-n-hexylbenzyl group, a 4-nonylbenzyl group, and a 3,4-dimethylbenzyl group.

12. (previously canceled)

13. (Currently Amended) A bathophenanthroline compound adapted to be used for use as an organic layer having a luminescent region provided between an anode and a cathode, wherein the organic layer comprises a bathophenanthroline compound of formula:

wherein Ar<sup>1</sup> and Ar<sup>2</sup> may be the same or different and independently represent an aryl group but do not form an interlocking macrocyclic compound.

EJ

14. (Currently Amended) The bathophenanthroline compound according to claim 13 wherein Ar¹ and Ar² are selected from the group consisting of a 1-naphthyl group, a 2-anthryl group, a 9-anthryl group, a 2-fluorenyl group, a 4-quinolyl group, a pyridyl group, a 3-pyridynyl group, a 2-pyridynyl group, a 2-furyl group, a 3-thienyl group, a 2-oxazolyl group, a 2-thiazolyl group, a 2-benzoxazoryl group, a 2-benzothiazoryl group, a 2-benzoimidazoryl group, a 4-methyphenyl group, a 3-methyphenyl group, a 3-methyphenyl group, a 2-methyphenyl group, a 3-methyphenyl n.n-dimethylphenyl group, a n.n-dimethylphenyl group, a n.n-dimethylphenyl group, a n.n-diethylphenyl group, a n-isopropylphenyl group, a 4-n-butylphenyl group, a 4-isobutylphenyl group, a n-tert-butylphenyl group.

- 15. (previously canceled)
- 16. (Previously Added) A process, comprising:
  - (a) obtaining a bathophenanthroline of formula

(b) subjecting the bathophenanthroline to nucleophilic substitution reaction at the 2, 9 positions by a lithium compound of formula (III):

R<sup>1</sup>-Li or R<sup>2</sup>-Li


wherein  $R^1$  and  $R^2$  may be the same or different and independently represent a hydrocarbon group provided that at least one of  $R^1$  and  $R^2$  has at least two carbon atoms, and  $R^1$  reacts at the 2 position and  $R^2$  reacts at the 9 position of the bathophenanthroline.

- 17. (Previously Added) The process according to Claim 16, wherein a carbanion is generated from the lithium compound in a solution and reacted with the bathophenanthroline during the nucleophilic substitution reaction.
- 18. (Previously Added) A process, comprising:
  - (a) obtaining a lithium compound of formula (V):

Ar<sup>1</sup>-Li or Ar<sup>2</sup>-Li

wherein Ar<sup>1</sup> and Ar<sup>2</sup> may be the same or different and independently represent an aryl group,

(b) subjecting the lithium compound to a bathophenanthroline of formula (IV):



via nucleophilic substitution reaction at the 2, 9 positions of the bathophenanthroline where  $A^1$  reacts at the 2 position and  $A^2$  reacts at the 9 position of the bathophenanthroline.

19. (Previously Added) The process according to Claim 18, wherein a carbanion is generated from the lithium compound in a solution and reacted with the bathophenanthroline during the nucleophilic substitution reaction.